

5314 DETROIT AVENUE LAKEWOOD, OH 44107

PHARMACY RH

PHARMACY DEPT NO. - 58330

RH:10698777666789 5V

STEVENS, STUART FRI

Mig: CIPROFLOXACIN

TAKE ONE OR TWO TABLETS BY MOUTH EVERY DAY

30 - 10 MG TABLETS

REFILLS: 2

UTR DR. L. ONYEJAMA, M.D. FAIRVIEW HOSPITAL

TELEPHONE: 216-222-2222

NO. 031

WARNINGS:
Do not take ciprofloxacin with dairy products such as milk or yogurt, or with calcium-fortified juice. Avoid taking antacids, iron, zinc, or aluminum supplements, tetracycline (Cephadril), or didanosine (didan) powder or chewable tablets within 2 hours before or after you take ciprofloxacin. Ciprofloxacin may cause swelling or tearing of a tendon (the thick cord that connects bones to muscles in the body), especially in the Achilles' tendon. You should not use ciprofloxacin if you are taking zanamivir (Zanamivir), if you have a history of myasthenia gravis, or if you are allergic to ciprofloxacin (Cipro) or similar antibiotics such as gemifloxacin (Factiva), levofloxacin (Levaquin), moxifloxacin (Avelox), norfloxacin (Noroxin), and others.
Before taking ciprofloxacin, tell your doctor if you have a heart rhythm disorder, kidney or liver disease, joint problems, diabetes, muscle weakness or trouble breathing, a history of seizures, low levels of potassium in your blood, a recent or family history of long QT syndrome, or if you have ever had an allergic reaction to an antibiotic.

5314 DETROIT AVENUE LAKEWOOD, OH 44107

PHARMACY RH

PHARMACY DEPT NO. - 58330

RH:10698777666789 5V

STEVENS, STUART FRI

Mig: CIPROFLOXACIN

TAKE ONE OR TWO TABLETS BY MOUTH EVERY DAY

30 - 10 MG TABLETS

REFILLS: 2

UTR DR. L. ONYEJAMA, M.D. FAIRVIEW HOSPITAL

TELEPHONE: 216-222-2222

NO. 031

WARNINGS:
Do not take ciprofloxacin with dairy products such as milk or yogurt, or with calcium-fortified juice. Avoid taking antacids, iron, zinc, or aluminum supplements, tetracycline (Cephadril), or didanosine (didan) powder or chewable tablets within 2 hours before or after you take ciprofloxacin. Ciprofloxacin may cause swelling or tearing of a tendon (the thick cord that connects bones to muscles in the body), especially in the Achilles' tendon. You should not use ciprofloxacin if you are taking zanamivir (Zanamivir), if you have a history of myasthenia gravis, or if you are allergic to ciprofloxacin (Cipro) or similar antibiotics such as gemifloxacin (Factiva), levofloxacin (Levaquin), moxifloxacin (Avelox), norfloxacin (Noroxin), and others.
Before taking ciprofloxacin, tell your doctor if you have a heart rhythm disorder, kidney or liver disease, joint problems, diabetes, muscle weakness or trouble breathing, a history of seizures, low levels of potassium in your blood, a recent or family history of long QT syndrome, or if you have ever had an allergic reaction to an antibiotic.

5314 DETROIT AVENUE LAKEWOOD, OH 44107

PHARMACY RH

PHARMACY DEPT NO. - 58330

RH:10698777666789 5V

STEVENS, STUART FRI

Mig: CIPROFLOXACIN

TAKE ONE OR TWO TABLETS BY MOUTH EVERY DAY

30 - 10 MG TABLETS

REFILLS: 2

UTR DR. L. ONYEJAMA, M.D. FAIRVIEW HOSPITAL

TELEPHONE: 216-222-2222

NO. 032

WARNINGS:
Do not take ciprofloxacin with dairy products such as milk or yogurt, or with calcium-fortified juice. Avoid taking antacids, iron, zinc, or aluminum supplements, tetracycline (Cephadril), or didanosine (didan) powder or chewable tablets within 2 hours before or after you take ciprofloxacin. Ciprofloxacin may cause swelling or tearing of a tendon (the thick cord that connects bones to muscles in the body), especially in the Achilles' tendon. You should not use ciprofloxacin if you are taking zanamivir (Zanamivir), if you have a history of myasthenia gravis, or if you are allergic to ciprofloxacin (Cipro) or similar antibiotics such as gemifloxacin (Factiva), levofloxacin (Levaquin), moxifloxacin (Avelox), norfloxacin (Noroxin), and others.
Before taking ciprofloxacin, tell your doctor if you have a heart rhythm disorder, kidney or liver disease, joint problems, diabetes, muscle weakness or trouble breathing, a history of seizures, low levels of potassium in your blood, a recent or family history of long QT syndrome, or if you have ever had an allergic reaction to an antibiotic.

5314 DETROIT AVENUE LAKEWOOD, OH 44107

PHARMACY RH

PHARMACY DEPT NO. - 58330

RH:10698777666789 5V

STEVENS, STUART FRI

Mig: CIPROFLOXACIN

TAKE ONE OR TWO TABLETS BY MOUTH EVERY DAY

30 - 10 MG TABLETS

REFILLS: 2

UTR DR. L. ONYEJAMA, M.D. FAIRVIEW HOSPITAL

TELEPHONE: 216-222-2222

NO. 032

WARNINGS:
Do not take ciprofloxacin with dairy products such as milk or yogurt, or with calcium-fortified juice. Avoid taking antacids, iron, zinc, or aluminum supplements, tetracycline (Cephadril), or didanosine (didan) powder or chewable tablets within 2 hours before or after you take ciprofloxacin. Ciprofloxacin may cause swelling or tearing of a tendon (the thick cord that connects bones to muscles in the body), especially in the Achilles' tendon. You should not use ciprofloxacin if you are taking zanamivir (Zanamivir), if you have a history of myasthenia gravis, or if you are allergic to ciprofloxacin (Cipro) or similar antibiotics such as gemifloxacin (Factiva), levofloxacin (Levaquin), moxifloxacin (Avelox), norfloxacin (Noroxin), and others.
Before taking ciprofloxacin, tell your doctor if you have a heart rhythm disorder, kidney or liver disease, joint problems, diabetes, muscle weakness or trouble breathing, a history of seizures, low levels of potassium in your blood, a recent or family history of long QT syndrome, or if you have ever had an allergic reaction to an antibiotic.

5314 DETROIT AVENUE LAKEWOOD, OH 44107

PHARMACY RH

PHARMACY DEPT NO. - 58330

RH:10698777666789 5V

STEVENS, STUART FRI

Mig: CIPROFLOXACIN

TAKE ONE OR TWO TABLETS BY MOUTH EVERY DAY

30 - 10 MG TABLETS

REFILLS: 2

UTR DR. L. ONYEJAMA, M.D. FAIRVIEW HOSPITAL

TELEPHONE: 216-222-2222

NO. 032

WARNINGS:
Do not take ciprofloxacin with dairy products such as milk or yogurt, or with calcium-fortified juice. Avoid taking antacids, iron, zinc, or aluminum supplements, tetracycline (Cephadril), or didanosine (didan) powder or chewable tablets within 2 hours before or after you take ciprofloxacin. Ciprofloxacin may cause swelling or tearing of a tendon (the thick cord that connects bones to muscles in the body), especially in the Achilles' tendon. You should not use ciprofloxacin if you are taking zanamivir (Zanamivir), if you have a history of myasthenia gravis, or if you are allergic to ciprofloxacin (Cipro) or similar antibiotics such as gemifloxacin (Factiva), levofloxacin (Levaquin), moxifloxacin (Avelox), norfloxacin (Noroxin), and others.
Before taking ciprofloxacin, tell your doctor if you have a heart rhythm disorder, kidney or liver disease, joint problems, diabetes, muscle weakness or trouble breathing, a history of seizures, low levels of potassium in your blood, a recent or family history of long QT syndrome, or if you have ever had an allergic reaction to an antibiotic.