

Directions

Use close cup, teaspoon (tsp), or tablespoon (TBSP)

Do not exceed 4 doses per 24 hours

Adults and children 12 years and over	1 TBSP (15 ml) every 6 - 8 hours
--	-------------------------------------

Children 6 to under 12 years	1 tsp (7 1.5 ml) every 6 - 8 hours
---------------------------------	---------------------------------------

Children under 6 years	Ask a doctor
---------------------------	--------------

www.vicks.com

Other Information

Each tablespoon contains sodium
28 mg. Store at room temperature

Inactive ingredients

Alcohol, citrix acid, FD&C Blue
No.1, FD&C Red No.40, flavour,
high fructose corn syrup, polyenth-
lene oxide, polyoxyl 40 stearate,
propylene glycol, purified water, ,
sodium citrate, sodium benzoate,
saccharin sodium.

Dist. by Procter & Gamble

VICKS

120 ml

**COUGH
SYRUP**

with
honey

**For dry and
irritating cough**



Drug Facts

Active Ingredient
(in each 15ml tablespoon)
Dextromethorphan H Br 30mg

Purpose
Cough Suppressant

Use
Temporarily relieves cough due
to minor throat and bronchial
irritation associated with a cold

Warnings

Do not use if you are now taking a
prescription monoamineoxidase
inhibitor (MAOI) (certain
drug for depression, psychiatric or
emotional conditions, or Parkison's
disease), or for 2 weeks after
stopping the MAOI drug. If you do not
know prescription drug contains an
MAIO, ask a doctor or pharmacist
before taking this product.

- Ask a doctor before use if you have
cough that occurs with too much
phlegm (mucus) persistent or chronic
cough such as occurs with smoking,
or emphysema

- Stop use and ask a doctor if cough
lasts more than 7 days, comes back,
or occurs with fever, rash, or
headache that lasts. These cough be
signs of a serious condition.

- If pregnant or breast-feeding, ask
a health professional before use.

VICKS

120 ml

**COUGH
SYRUP**

with
honey

**For dry and
irritating cough**

